

GET READY TO EXPLORE NEW POSSIBILITIES WITH QFITLIA[™]

This guide can help you organize your thoughts and make the most of your discussion with your doctor to discover if Qfitlia is right for you

INDICATION

Qfitlia[™] (fitusiran) is a prescription medicine used for routine prophylaxis to prevent or reduce the frequency of bleeding episodes in adults and children 12 years and older with hemophilia A or B with or without Factor VIII or IX inhibitors.

It is not known if Qfitlia is safe and effective in children younger than 12 years of age.

IMPORTANT SAFETY INFORMATION

Qfitlia can cause SERIOUS SIDE EFFECTS, including:

- **Abnormal blood clotting (thrombotic events):** Serious blood clots have occurred in people treated with Qfitlia. Qfitlia can cause blood clots to form in the blood vessels in your arms, legs, lungs, heart, brain, eyes, or head. Your risk of blood clots is greater if your antithrombin (AT) blood level is persistently less than 15% or if you have certain other conditions. Your healthcare provider (HCP) will check your AT blood levels before and during treatment with Qfitlia
- **Gallbladder disease:** Qfitlia can cause gallstones and inflammation of your gallbladder, which might require surgery to remove your gallbladder. Tell your HCP right away if you develop stomach pain, indigestion, nausea, or vomiting. Your HCP may temporarily or permanently stop Qfitlia if you develop any of these symptoms

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#), including SERIOUS SIDE EFFECTS, and [Medication Guide](#).


Reflect on how your current hemophilia management supports your lifestyle

How often do you administer your current treatment?

Do you find it easy or difficult to stick to this treatment plan?

How many bleeds did you have in the past year?

To what extent have you limited planning activities while on your current treatment?



What are your hemophilia treatment goals?

(e.g., protection from bleeds, lifestyle suitability, convenience of administration, travel-friendly)

Speak with your doctor to discover more about Qfitlia™

Questions to get you started:

- | | |
|---|--|
| <input type="checkbox"/> What are the potential benefits and side effects of Qfitlia? | <input type="checkbox"/> Are there any tests I need to take with Qfitlia and how often do I need to take them? |
| <input type="checkbox"/> What should I do if I get a bleed while taking Qfitlia? | <input type="checkbox"/> How does Qfitlia work? |
| <input type="checkbox"/> What should I know about starting treatment with Qfitlia? | <input type="checkbox"/> Is Qfitlia right for me? |
| <input type="checkbox"/> What dose of Qfitlia will I need and how often will I need to take it? | |

Additional questions/notes:

IMPORTANT SAFETY INFORMATION (CONT'D)

What is the most important information I should know about Qfitlia?

Qfitlia helps your blood form clots. Do not stop using Qfitlia without talking to your HCP. If you miss doses or stop using Qfitlia, you may no longer be protected against bleeding.

Use of a clotting factor concentrate (CFC) or bypassing agent (BPA) to help protect against bleeding must be stopped within 7 days after your first dose of Qfitlia.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#), including **SERIOUS SIDE EFFECTS, and [Medication Guide](#).**

6 THINGS TO KNOW ABOUT QFITLIA™

Qfitlia is an antithrombin-lowering agent for people 12 years and older with hemophilia A or B, with or without inhibitors



Qfitlia prophylaxis helps your blood's ability to clot by lowering a protein called antithrombin



Consistent protection with as few as 6 injections a year*



Easy-to-use prefilled pen†



No refrigeration for up to 3 months‡



No mixing needed



No weight-based dosing



Speak with your doctor to find out more



Looking for more information?

HemAssist™ Patient Support from Sanofi supports you with treatment access, financial assistance, and education throughout your journey with Qfitlia

[Learn more](#)

* In the clinical study of 227 people, ~67% of people with hemophilia took Qfitlia every other month, and ~19% took it monthly.

† Prefilled pen administration for the 50 mg dose. Vial and syringe administration for the 10 mg and 20 mg dose.

‡ Prefilled pen may be stored at room temperature between 59 °F to 86 °F (15 °C to 30 °C) for a single period of up to 3 months within the expiration date printed on the label.

IMPORTANT SAFETY INFORMATION (CONT'D)

What is the most important information I should know about Qfitlia? (cont'd)

Your HCP may prescribe on-demand CFC or BPA if you bleed during treatment with Qfitlia. **Carefully follow your HCP's instructions regarding when to use on-demand treatment with CFC or BPA, including the prescribed dose and timing of the CFC or BPA.**

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#), including **SERIOUS SIDE EFFECTS**, and [Medication Guide](#).

Qfitlia™
(fitusiran) injection
50 mg/0.5 mL

INDICATION

Qfitlia™ (fitusiran) is a prescription medicine used for routine prophylaxis to prevent or reduce the frequency of bleeding episodes in adults and children 12 years and older with hemophilia A or B with or without Factor VIII or IX inhibitors.

It is not known if Qfitlia is safe and effective in children younger than 12 years of age.

IMPORTANT SAFETY INFORMATION

Qfitlia can cause SERIOUS SIDE EFFECTS, including:

- **Abnormal blood clotting (thrombotic events):** Serious blood clots have occurred in people treated with Qfitlia. Qfitlia can cause blood clots to form in the blood vessels in your arms, legs, lungs, heart, brain, eyes, or head. Your risk of blood clots is greater if your antithrombin (AT) blood level is persistently less than 15% or if you have certain other conditions. Your healthcare provider (HCP) will check your AT blood levels before and during treatment with Qfitlia
- **Gallbladder disease:** Qfitlia can cause gallstones and inflammation of your gallbladder, which might require surgery to remove your gallbladder. Tell your HCP right away if you develop stomach pain, indigestion, nausea, or vomiting. Your HCP may temporarily or permanently stop Qfitlia if you develop any of these symptoms

What is the most important information I should know about Qfitlia?

Qfitlia helps your blood form clots. Do not stop using Qfitlia without talking to your HCP. If you miss doses or stop using Qfitlia, you may no longer be protected against bleeding.

Use of a clotting factor concentrate (CFC) or bypassing agent (BPA) to help protect against bleeding must be stopped within 7 days after your first dose of Qfitlia.

Your HCP may prescribe on-demand CFC or BPA if you bleed during treatment with Qfitlia. **Carefully follow your HCP's instructions regarding when to use on-demand treatment with CFC or BPA, including the prescribed dose and timing of the CFC or BPA.**

Get medical help right away if you get any of these signs or symptoms of blood clots during or after treatment with Qfitlia:

- Swelling, pain, or redness in arms or legs
- Coughing up blood
- Shortness of breath
- Severe chest pain or tightness of the chest
- Fast heart rate
- Feeling faint or passing out
- Severe or persistent headache
- Difficulty speaking or understanding language
- Feeling confused
- Numbness or weakness in your face, arms, or legs
- Sudden loss or changes in your vision, eye pain, or swelling

What are the possible side effects of Qfitlia?

- **Qfitlia can cause other serious side effects, including an increase in your blood liver enzymes.** Your HCP will do blood tests to check your liver function before and during treatment with Qfitlia
- **The most common side effects of Qfitlia include** viral infection, common cold symptoms, and bacterial infection

These are not all the possible side effects of Qfitlia.

What should I tell my HCP before using Qfitlia?

- **Tell your HCP about all of your medical conditions,** including if you have liver problems, a history of gallbladder disease, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed
- **Females who are able to become pregnant:** Hormonal birth control may increase your risk of blood clots if used during treatment with Qfitlia. Talk to your HCP about effective forms of non-hormonal birth control you can use before starting and during treatment with Qfitlia
- **Tell your HCP about all of the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements

Please see full [Prescribing Information](#), including **SERIOUS SIDE EFFECTS**, and [Medication Guide](#).